

# MAX Function

Computes the maximum value found in all row values in a column. Input column are permitted to be of Integer or Decimal.

- When used in a `pivot` transform, the function is computed for each instance of the value specified in the `group` parameter. See *Pivot Transform*.
- If a row contains a missing or null value, it is not factored into the calculation.
- If no numeric values are found in the source column, the function returns a null value.

For a version of this function computed over a rolling window of rows, see *ROLLINGMAX Function*.

## Basic Usage

```
pivot value:MAX(myRating) group:postal_code limit:1
```

**Output:** Generates a two-column table containing the unique values from the `postal_code` column and the maximum value of the `myRating` column for the `postal_code` value . The `limit` parameter defines the maximum number of output columns.

## Syntax

```
pivot value:MAX(function_col_ref) [group:group_col_ref] [limit:limit_count]
```

Argument	Required?	Data Type	Description
<code>function_col_ref</code>	Y	string	Name of column to which to apply the function

For more information on the `group` and `limit` parameters, see *Pivot Transform*.

For more information on syntax standards, see *Language Documentation Syntax Notes*.

## function\_col\_ref

Name of the column the values of which you want to calculate the maximum. Column must contain Integer or Decimal values.

- Literal values are not supported as inputs.
- Multiple columns and wildcards are not supported.

### Usage Notes:

Required?	Data Type	Example Value
Yes	String (column reference)	<code>myValues</code>

## Examples

This example illustrates how you can apply statistical functions to your dataset. Calculations include average (mean), max, min, standard deviation, and variance.

### Source:

Students took a test and recorded the following scores. You want to perform some statistical analysis on them:

Student	Score
Anna	84
Ben	71
Caleb	76
Danielle	87
Evan	85
Faith	92
Gabe	85
Hannah	99
Ian	73
Jane	68

**Transform:**

You can use the following transforms to calculate the average (mean), minimum, and maximum scores:

```
derive type:single value:AVERAGE(Score) as:'avgScore'
```

```
derive type:single value:MIN(Score) as:'minScore'
```

```
derive type:single value:MAX(Score) as:'maxScore'
```

To apply statistical functions to your data, you can use the VAR and STDEV functions, which can be used as the basis for other statistical calculations.

```
derive type:single value:VAR(Score)
```

```
derive type:single value:STDEV(Score)
```

For each score, you can now calculate the variation of each one from the average, using the following:

```
derive type:single value:((Score - avg_Score) / stdev_Score) as:'stDevs'
```

Now, you want to apply grades based on a formula:

Grade	standard deviations from avg (stDevs)
A	stDevs > 1
B	stDevs > 0.5
C	-1 <= stDevs <= 0.5
D	stDevs < -1
F	stDevs < -2

You can build the following transform using the IF function to calculate grades.

```
derive type:single value:IF((stDevs > 1),'A',IF((stDevs < -2),'F',IF((stDevs < -1),'D',IF
((stDevs > 0.5),'B','C'))))
```

For more information, see *IF Function*.

To clean up the content, you might want to apply some formatting to the score columns. The following reformats the `stdev_Score` and `stDevs` columns to display two decimal places:

```
set col:stdev_Score value:NUMFORMAT(stdev_Score, '##.00')
```

```
set col:stDevs value:NUMFORMAT(stDevs, '##.00')
```

```
derive type:single value:MODE(Score) as:'modeScore'
```

### Results:

Student	Score	modeScore	avgScore	minScore	maxScore	var_Score	stdev_Score	stDevs	Grade
Anna	84	85	82	68	99	87.000000000000001	9.33	0.21	C
Ben	71	85	82	68	99	87.000000000000001	9.33	-1.18	D
Caleb	76	85	82	68	99	87.000000000000001	9.33	-0.64	C
Danielle	87	85	82	68	99	87.000000000000001	9.33	0.54	B
Evan	85	85	82	68	99	87.000000000000001	9.33	0.32	C
Faith	92	85	82	68	99	87.000000000000001	9.33	1.07	A
Gabe	85	85	82	68	99	87.000000000000001	9.33	0.32	C
Hannah	99	85	82	68	99	87.000000000000001	9.33	1.82	A
Ian	73	85	82	68	99	87.000000000000001	9.33	-0.96	C
Jane	68	85	82	68	99	87.000000000000001	9.33	-1.50	D